

ARTÍCULO DE MAYOR FACTOR DE IMPACTO MAYO 2013

Fernandez-Hidalgo, N; Almirante, B; Gavalda, J; Gurgui, M; Pena, C; de Alarcon, A; Ruiz, J; **Vilacosta, I**; Montejo, M; Vallejo, N; Lopez-Medrano, F; Plata, A; Lopez, J; Hidalgo-Tenorio, C; Galvez, J; Saez, C; Lomas, JM; Falcone, M; de la Torre, J; Martinez-Lacasa, X; Pahissa, A. Ampicillin Plus Ceftriaxone Is as Effective as Ampicillin Plus Gentamicin for Treating *Enterococcus faecalis* Infective Endocarditis. CLINICAL INFECTIOUS DISEASES 2013. 56 (9):1261 - 1268. doi: 10.1093/cid/cit052. FACTOR DE IMPACTO: **9,154**.


OXFORD JOURNALS

Clinical Infectious Diseases

ABOUT THIS JOURNAL CONTACT THIS JOURNAL SUBSCRIPTIONS CURR

Oxford Journals > Medicine > Clinical Infectious Diseases > Volume 56, Issue 9 > Pp. 1261-1268.

Stay up to date in Infectious Diseases



Ampicillin Plus Ceftriaxone Is as Effective as Ampicillin Plus Gentamicin for Treating *Enterococcus faecalis* Infective Endocarditis

Nuria Fernández-Hidalgo¹, Benito Almirante¹, Joan Gavalda¹, Mercè Gurgui², Carmen Peña³, Aristides de Alarcón⁴, Josefa Ruiz⁵, Isidre Vilacosta⁶, Miguel Montejo⁷, Nuria Vallejo⁸, Francisco López-Medrano⁹, Antonio Plata¹⁰, Javier López¹¹, Carmen Hidalgo-Tenorio¹², Juan Gálvez¹³, Carmen Sáez¹⁴, José Manuel Lomas¹⁵, Marco Falcone¹⁶, Javier de la Torre¹⁶, Xavier Martínez-Lacasa¹⁷, and Albert Pahissa¹

[Author Affiliations](#)

Correspondence: Nuria Fernández-Hidalgo, MD, Servei de Malalties Infeccioses, Hospital Universitari Vall d'Hebron, Passeig de la Vall d'Hebron 119-129, 08035 Barcelona, Spain (nufernan@gmail.com).

Abstract

Background. The aim of this study was to compare the effectiveness of the ampicillin plus ceftriaxone (AC) and ampicillin plus gentamicin (AG) combinations for treating *Enterococcus faecalis* infective endocarditis (EFIE).